arXiv:1303.3114v1 [math.FA] 13 Mar 2013

A NOTE ON SANTALÓ INEQUALITY FOR THE POLARITY TRANSFORM AND ITS REVERSE

SHIRI ARTSTEIN-AVIDAN AND BOAZ A. SLOMKA

Abstract. We prove a Santaló and a reverse Santaló inequality for the class consisting of even log-concave functions attaining their maximal value 1 at the origin, also called even geometric log-concave functions. We prove that there exist universal numerical constants $c, C > 0$ such that for any even geometric log-concave function $f = e^{-\varphi}$,

$$c^n \cdot |B_2^n|^2 \le \int_{\mathbb{R}^n} e^{-\varphi} \int_{\mathbb{R}^n} e^{-\varphi^\circ} \le C^n \cdot n!^2 \, |B_2^n|^2$$

where B_2^n is the Euclidean unit ball of $\mathbb{R}^n$ and φ° is the polar function of φ (not the Legendre transform!), a transform which was recently re-discovered by Artstein-Avidan and Milman, and is defined below. The bounds are sharp up to the optimal constants c, C.

1. Introduction and main results

Let $\mathbb{R}^n$ denote the Euclidean n-dimensional real space, equipped with the standard scalar product $\langle \cdot, \cdot \rangle$ and the standard Euclidean norm $|x| = \langle x, x \rangle^{1/2}$. Denote its unit ball by B_2^n. Let $K \subseteq \mathbb{R}^n$ be a centrally symmetric convex body (i.e., compact and containing the origin 0 in its interior). The polar set of K is given by

$$K^\circ = \left\{ x \in \mathbb{R} \,:\, \sup_{y \in K} \langle x, y \rangle \le 1 \right\}$$

and is also a centrally symmetric convex body. Denote the Lebesgue volume of K by $|K|$. Recall the classical Santaló inequality and its reverse: there exists an absolute constant $c > 0$ such that for any dimension n and every centrally symmetric convex body $K \subseteq \mathbb{R}^n$,

$$c^n \cdot |B_2^n|^2 \le |K| \, |K^\circ| \le |B_2^n|^2 \, .$$

The right hand side is referred to as Santaló inequality and is due to Santaló [16] (for a simpler proof see [11]). The left hand side inequality is referred to as the reverse Santaló inequality, or the Bourgain-Milman inequality and is due to J. Bourgain and V. Milman [5]. While bounds for the numerical constant c have been improving over the years (see [8, 13]) it is still an open question whether among centrally symmetric convex bodies the unit cube minimizes this product, often called the Mahler product.

Functional versions of Santaló inequality and its reverse were also established; A function $f : \mathbb{R}^n \to \mathbb{R}$ is said to be log-concave if it is of the form $e^{-\varphi}$ where $\varphi : \mathbb{R}^n \to \mathbb{R} \cup \{\infty\}$ is convex. One interpretation for the "polar" function of $e^{-\varphi}$ may be the function $e^{-\mathcal{L}\varphi}$ where $\mathcal{L}$ is the well-known Legendre transform, given by $\mathcal{L}\varphi(x) = \sup_{y \in \mathbb{R}^n} [\langle x, y \rangle - \varphi(y)]$. Note that $e^{-\mathcal{L}\varphi}$ is always log-concave. It was proven that there exists an absolute constant $c > 0$ such that for any even log-concave function $f = e^{-\varphi}$ on $\mathbb{R}^n$ with $0 < \int f < \infty$,

$$\left(\frac{2\pi}{c} \right)^n \le \int_{\mathbb{R}^n} e^{-\varphi} \cdot \int_{\mathbb{R}^n} e^{-\mathcal{L}\varphi} \le (2\pi)^n \, .$$

2010 *Mathematics Subject Classification.* 52A41; 26A51; 46B10.

Key words and phrases. Santaló and reverse Santaló inequality, polarity transform, log-concave function.

This work was supported by ISF¸grant No. 247/11.

The right hand side is due to Ball [4], see also Artstein, Klartag, and Milman [1]. In the latter paper it is actually proved that Gaussians are the only maximizers for this product. For further related results see also [6, 9, 10]. The left hand side is due to Klartag and Milman [7]. Note that Santaló inequality and its reverse for convex bodies may be recovered by plugging in $f = e^{-\|\cdot\|_K^2/2}$, where $\|\cdot\|_K$ denotes the norm with unit ball K.

In this note, we prove a different functional version of Santaló inequality and its reverse, for an important subclass of even log-concave functions, called *(even) geometric log-concave functions*, and consisting of all functions of the form $f = e^{-\varphi}$ where φ is a non-negative even lower semi-continuous convex function with $\varphi(0) = 0$. A few years ago, Artstein-Avidan and Milman [2, 3] have proven that, on this class, up to trivial obvious modifications, there exist exactly two order-reversing involutions, one is the Legendre transform, and the other, which we shall call here the polarity transform, is defined by $\varphi \mapsto \varphi^\circ$ where

$$\varphi^\circ(x) = \sup_{y\in\mathbb{R}^n} \frac{\langle x, y\rangle - 1}{\varphi(y)}.$$

Here we agree that $\frac{+}{0} = \infty$, $\frac{0}{0} = 0$ and $\frac{-}{0} = \left(\frac{-}{0}\right)_+ = 0$. The latter is needed only for this definition to work with the identically 0 function. It seems that this transform has appeared only once in the literature (Rockafeller's book [14, p. 136]) before it was re-discovered in [2, 3]. Among other things, in the latter paper strong reasons were given to explain why this new transform should be considered as the natural extension of the notion of polarity from convex bodies to the class of geometric log-concave functions, and the Legendre transform as the natural extension of the support function.

Since these discoveries, the first named author was asked in several occasions whether a Santaló type inequality holds also for the polarity transform. In this note we answer this question in the affirmative. We prove the following:

Theorem 1. *Let $f = e^{-\varphi}$ be an even geometric log-concave function on $\mathbb{R}^n$ such that $0 < \int e^{-\varphi} < \infty$. Then*

$$c^n \cdot |B_2^n|^2 \le \int_{\mathbb{R}^n} e^{-\varphi} \cdot \int_{\mathbb{R}^n} e^{-\varphi^\circ} \le C^n \cdot n!^2 \, |B_2^n|^2$$

where $c, C > 0$ are universal constants independent of n and φ.

Note that, up to optimal constants c and C, one cannot hope for better bounds since for $\varphi = e^{-|x|}$ the above product equals $n!^2 \cdot |B_2^n|^2$ and for indicators $\mathbb{1}_K$ of centrally symmetric convex bodies $K \subseteq \mathbb{R}^n$ we recover the Bourgain-Milman lower bound. However, although the product $\int \varphi \int \varphi^\circ$ is invariant under invertible linear transformations, as one can check that $(\varphi \circ A)^\circ = \varphi^\circ \circ A^{-T}$, it does not imply that this product has maximizers or minimizers. We remark also that the fixed points of the mapping $\varphi \mapsto \varphi^\circ$ in dimension 1 have been classified by L. Rotem in [15]. The motivation for classifying these fixed points was that in the convex body case the maximizer for the Mahler product is the unique fixed point of the polarity transform for convex bodies, the Euclidean ball. Thus the maximizer of the product considered in Theorem 1, if exists, might also be a fixed point, although this was not verified. We do know to show, by symmetrizations, that maximizers, if exist, must be rotationally invariant.

This note is organized as follows. In Section 2.1 we prove some preliminary facts about log-concave functions, in particular that the class of geometric log-concave functions with (finite) positive integral is closed under the polarity transform, and a proposition that appeared in [12] about the connection between level sets of a function and its polar, to which we give an alternative geometric proof. The proof of Theorem 1 is split into two propositions, the right hand side inequality is proven in Section 3 and the left hand side inequality is proven in Section 4.

2. PRELIMINARY FACTS

2.1. **Comparing level sets of a function and its polar.** Let $f = e^{-\varphi}$ be a geometric log-concave function on $\mathbb{R}^n$. We adopt the notation of [12] and denote the level sets of f by

$$\overline{K}_t\left(f\right) := \left\{x \in \mathbb{R}^n \,:\, f\left(x\right) \geq t\right\}$$

for any $0 < t \leq 1$. Similarly we denote the level sets of φ by

$$\underline{K}_t\left(\varphi\right) := \left\{x \in \mathbb{R}^n \,:\, \varphi\left(x\right) \leq t\right\}$$

for any $0 \leq t < \infty$. Note that $\overline{K}_t\left(f\right)$ is a closed convex set, and also that $\overline{K}_t\left(f\right) = \underline{K}_{\ln(1/t)}\left(\varphi\right)$. Moreover, note that if f is even then $\overline{K}_t\left(f\right)$ are centrally symmetric. V. Milman and L. Rotem [12, Proposition 11] proved the following useful proposition:

Proposition 2. *For any geometric convex function* $\varphi : \mathbb{R}^n \to [0, \infty]$ *we have that*

$$\left(\underline{K}_{1/t}\left(\varphi\right)\right)^\circ \subseteq \underline{K}_t\left(\varphi^\circ\right) \subseteq 2\left(\underline{K}_{1/t}\left(\varphi\right)\right)^\circ$$

For the reader's convenience, we provide an alternative geometric proof to the above proposition. To this end, we need to recall the following simple facts about the polarity transform (for proofs see e.g., [3]):

Facts. For any geometric convex functions φ, ψ on $\mathbb{R}^n$ we have that

1. $\varphi \leq \psi$ if and only if $\psi^\circ \leq \varphi^\circ$. Also $\left(\varphi^\circ\right)^\circ = \varphi$.
2. $\max\left(\varphi, \psi\right) = \hat{\min}\left(\varphi^\circ, \psi^\circ\right)$ where $\hat{\min}\left(\varphi, \psi\right) = \sup\left\{\phi \text{ a geometric convex function } : \phi \leq \varphi \text{ and } \phi \leq \psi\right\}$.
3. For any centrally symmetric convex body K and any $t > 0$, $\left(t\left\|\cdot\right\|_K\right)^\circ = \frac{1}{t}\left\|\cdot\right\|_{K^\circ}$ and $\left(\mathbb{1}_K^\infty\right)^\circ = \mathbb{1}_{K^\circ}^\infty$, where $\mathbb{1}_K^\infty = -\log\left(\mathbb{1}_K\right)$.

Proof of Proposition 2. Let $t > 0$. Since φ is a geometric convex function, it readily follows that for all $x \in \underline{K}_t\left(\varphi\right)$,

$$\mathbb{1}_{\underline{K}_t}^\infty\left(x\right) \leq \varphi\left(x\right) \leq t \cdot \left\|x\right\|_{\underline{K}_t(\varphi)}$$

and for all $x \notin \underline{K}_t\left(\varphi\right)$,

$$t \cdot \left\|x\right\|_{\underline{K}_t(\varphi)} \leq \varphi\left(x\right) \leq \mathbb{1}_{\underline{K}_t}^\infty\left(x\right).$$

Thus, the following inequality holds:

$$\hat{\min}\left(t \cdot \left\|\cdot\right\|_{\underline{K}_t(\varphi)}, \mathbb{1}_{\underline{K}_t(\varphi)}^\infty\right) \leq \varphi \leq \max\left(t \cdot \left\|\cdot\right\|_{\underline{K}_t(\varphi)}, \mathbb{1}_{\underline{K}_t(\varphi)}^\infty\right).$$

Applying the inequality with the polar transform and using Facts 1-3 yields

$$\psi_1 := \hat{\min}\left(\frac{1}{t} \cdot \left\|\cdot\right\|_{\underline{K}_t^\circ(\varphi)}, \mathbb{1}_{\underline{K}_t^\circ(\varphi)}^\infty\right) \leq \varphi^\circ \leq \max\left(\frac{1}{t} \cdot \left\|\cdot\right\|_{\underline{K}_t^\circ(\varphi)}, \mathbb{1}_{\underline{K}_t^\circ(\varphi)}^\infty\right) =: \psi_2.$$

As illustrated in Fig. 2.1, it is easy to check that $\underline{K}_{1/t}\left(\psi_2\right) = \underline{K}_t^\circ\left(\varphi\right)$, and so the right hand side inequality implies that $\underline{K}_t^\circ\left(\varphi\right) \subseteq K_{1/t}\left(\varphi^\circ\right)$. Similarly, $\underline{K}_{1/t}\left(\psi_1\right) = 2 \cdot \underline{K}_t^\circ\left(\varphi\right)$, and so the left hand side inequality implies that $\underline{K}_{1/t}\left(\varphi^\circ\right) \subseteq 2 \cdot \underline{K}_t^\circ\left(\varphi\right)$.

□


$\underline{K}_{1/t}(\varphi^\circ)$
$\underline{K}_{1/t}(\psi_2)$
ψ_2
φ°
$\underline{K}^\circ_t(\varphi)$
$\underline{K}_{1/t}(\psi_1)$
$\underline{K}_{1/t}(\varphi^\circ)$
φ°
ψ_1
$\underline{K}^\circ_t(\varphi)$

(A) $\underline{K}^\circ_t(\varphi) \subseteq K_{1/t}(\varphi^\circ)$ (B) $K_{1/t}(\varphi^\circ) \subseteq 2\cdot \underline{K}^\circ_t(\varphi)$

FIGURE 2.1. Comparing level sets of φ and φ° in Proposition 2

2.2. **Log-concave functions with finite positive integral.** In this section we examine some properties of geometric log-concave functions f satisfying that $0<\int f<\infty$. As later on we will be connecting the integral of such a function to an integral over its level sets, it will be helpful to show that its level sets are convex bodies, that is (closed) convex sets with a finite positive volume. This will allow us to apply the classical Santaló (or reverse Santaló) inequality for convex bodies in our setting. To this end, denote the support of a log-concave function f by $\operatorname{supp}(f)=\{x:\mathbb{R}^n : f(x)>0\}$. The following holds.

Lemma 3. *Let* $f=e^{-\varphi}$ *be a geometric log-concave function on* $\mathbb{R}^n$. *Then* $0<\int f<\infty$ *if and only if for some open ball* $B\subseteq\mathbb{R}^n$ *and* $0<\varepsilon<1$, $\varepsilon\cdot\mathbb{1}_B(x)\le f(x)$ *and for some* $r,c>0$, $f(x)\le e^{-c|x|}$ *for all* $|x|\ge r$.

Proof. Assume that $0<\int f<\infty$. The fact that $0<\int f$ obviously implies that $\operatorname{supp}(f)$ is of positive Lebesgue measure. Since $\operatorname{supp}(f)$ is a convex set, and f is continuous on its support, there exist $0<\varepsilon<e^{-1}$ and an open ball $B\subseteq\operatorname{supp}(f)$ such that $f(B)\ge\varepsilon$. Denote the unit sphere in $\mathbb{R}^n$ by $\mathbb{S}^{n-1}=\{\theta\in\mathbb{R}^n : |\theta|=1\}$. Then for each $\theta\in\mathbb{S}^{n-1}$ there exists $R>0$ such that $f(R\theta)\le e^{-1}$. Indeed, otherwise we would have that $f(\{\theta r : r\ge 0\})>e^{-1}$ for some $\theta\in\mathbb{S}^{n-1}$and so, by the log-concavity of f, $f\ge\varepsilon$ on the convex hull of B and the ray $\{\theta r : r\ge0\}$, which has infinite Lebesgue measure, and so $\int f=\infty$, a contradiction. Hence, by the compactness of $\mathbb{S}^{n-1}$ there exists $r=\min\left\{R>0 : f(\theta R)\le e^{-1}\text{for all } \theta\in\mathbb{S}^{n-1}\right\}$, and so for every $x\in\mathbb{R}^n$ with $|x|\ge r$,

$$e^{-1}\ge f\left(\frac{r}{|x|}x\right)=f\left(\frac{r}{|x|}x+\left(1-\frac{r}{|x|}\right)0\right)\ge f(x)^{r/|x|}f(0)^{1-r/|x|}=f(x)^{r/|x|}$$

from which it follows that $f\le e^{-|x|/r}$ for all $|x|\ge r$.

For the opposite direction we have:

$$0<\int\varepsilon\cdot\mathbb{1}_B<\int e^{-\varphi}<\int_{|x|<r}e^{-\varphi}+\int_{|x|\ge r}e^{-\varphi}\le|rB_2^n|+\int e^{-c|x|}<\infty$$

□

Remark 4. One may check that the above lemma is equivalent to the fact that a geometric log-concave function f has a finite positive integral if and only if its support is of full dimension and f does not attain the constant value 1 on a whole ray.

As a consequence of the above lemma we can conclude that the level sets of a geometric log-concave function $f = e^{-\varphi}$ satisfying that $0 < \int f < \infty$ are convex bodies, and that the class of such functions is closed under the polarity transform, that is $0 < \int e^{-\varphi^\circ} < \infty$:

Proposition 5. *Let $f = e^{-\varphi}$ be a geometric log-concave function on $\mathbb{R}^n$, satisfying that $0 < \int f < \infty$. Then for all $t > 0$, $\underline{K}_t(\varphi)$ and $\underline{K}_t(\varphi^\circ)$ are compact convex sets with non-vanishing Lebesgue volume. Moreover, $0 < \int e^{-\varphi^\circ} < \infty$.*

Proof. By Lemma 3, $f(x) \le e^{-c|x|}$ for all $|x| \ge r$ and $\varepsilon \cdot \mathbb{1}_B \le f$ for some $0 < \varepsilon < 1$ and an open ball B. Let $t > 0$ such that $x \in \overline{K}_t(f)$ for some $|x| \ge r$. Since $f(x) \le e^{-c|x|}$ for all $|x| \ge r$, it follows that $x \in \overline{K}_t\left(e^{-c|\cdot|}\right)$ and so $\left|\overline{K}_t(f)\right| \le \left|\overline{K}_t\left(e^{-c|\cdot|}\right)\right| < \infty$. By the same reasoning the last inequality holds for all $s < t$. As $\overline{K}_t(f)$ is monotonically decreasing with t, we have that $\left|\overline{K}_t(f)\right| < \infty$ for all $0 < t < 1$. In other words $|\underline{K}_t(\varphi)| < \infty$ for all $t > 0$. Since $\varepsilon \cdot \mathbb{1}_B \le f$ it follows that $\overline{K}_t(\varepsilon \cdot \mathbb{1}_B) \subseteq \overline{K}_t(f)$, and so $0 < \left|\overline{K}_t(\varepsilon \cdot \mathbb{1}_B)\right| \le \left|\overline{K}_t(f)\right|$, for all $0 < t \le \varepsilon$. In other words $0 < |\underline{K}_t(\varphi)|$ for all $a := \log(1/\varepsilon) \le t$. As φ is a geometric log-concave function, we have that $\varphi(x) \le a \cdot \|x\|_{\underline{K}_a(\varphi)}$ for all $x \in \underline{K}_a(\varphi)$ which means that $\underline{K}_t\left(a \cdot \|x\|_{\underline{K}_a(\varphi)}\right) \subseteq \underline{K}_t(\varphi)$ for all $0 < t < a$ and hence $0 < |\underline{K}_t(\varphi)|$ for all $0 < t$. Proposition 2 then implies that

$$0 < \left|\underline{K}_{1/t}(\varphi)\right| \le |\underline{K}_t(\varphi^\circ)| \le 2^n \cdot \left|\underline{K}_{1/t}(\varphi)\right| < \infty.$$

Denote $g(x) = e^{-\varphi^\circ(x)}$. Since $0 < |\underline{K}_1(\varphi^\circ)| < \infty$ and $\underline{K}_1(\varphi^\circ)$ is convex, there exist an open ball B and a radius $r > 0$ such that $B \subseteq \underline{K}_1(\varphi^\circ) \subseteq rB_2^n$, and so on the one hand

$$\frac{1}{e} \cdot \mathbb{1}_B \le \frac{1}{e} \cdot \mathbb{1}_{\underline{K}_1(\varphi^\circ)} = \frac{1}{e} \cdot \mathbb{1}_{\overline{K}_{1/e}(g)} \le g = e^{-\varphi^\circ}.$$

On the other hand, for some $c > 0$ we have that $\varphi^\circ(x) \ge \|x\|_{\underline{K}_1(\varphi^\circ)} \ge c\,|x|$ for all $x \notin \underline{K}_1(\varphi^\circ)$ and in particular for all $|x| \ge r$. In other words, for all $|x| \ge r$ we have that $e^{-\varphi^\circ} \le e^{-c|x|}$. Thus, Lemma 3 implies that $0 < \int e^{-\varphi^\circ} < \infty$. ☐

3. SANTALÓ INEQUALITY FOR THE POLARITY TRANSFORM

Proposition 6. *Let φ be a geometric even convex function on $\mathbb{R}^n$ with $0 < \int e^{-\varphi} < \infty$. Then*

$$\int_{\mathbb{R}^n} e^{-\varphi} \cdot \int_{\mathbb{R}^n} e^{-\varphi^\circ} \le C^n \cdot n!^2 \left|B_2^n\right|^2$$

where $C > 0$ is a universal constant, independent of n and φ.

Proof. We have that

$$\int_{\mathbb{R}^n} e^{-\varphi} = \int_0^1 \left|\overline{K}_t\left(e^{-\varphi}\right)\right| dt = \int_0^1 \left|\underline{K}_{\ln(1/t)}(\varphi)\right| dt = \int_0^\infty e^{-s} \cdot |\underline{K}_s(\varphi)|\, ds.$$

By the convexity of φ, we have that $\varphi(x) \ge \|x\|_{\underline{K}_1(\varphi)}$ for all $x \notin \underline{K}_1(\varphi)$, from which it follows that $\underline{K}_s(\varphi) \subseteq \underline{K}_s\left(\|\cdot\|_{\underline{K}_1(\varphi)}\right)$ for all $s > 1$. Thus, for some $c_1, c_2 > 0$,

$$\begin{aligned}
\int_0^\infty e^{-s} \cdot |\underline{K}_s(\varphi)|\, ds &= \int_0^1 e^{-s} \cdot |\underline{K}_s(\varphi)|\, ds + \int_1^\infty e^{-s} \cdot |\underline{K}_s(\varphi)|\, ds \\
&\le |\underline{K}_1(\varphi)| \cdot \int_0^1 e^{-s} ds + \int_1^\infty e^{-s} \cdot \left|\underline{K}_s\left(\|\cdot\|_{\underline{K}_1(\varphi)}\right)\right| ds \\
&= c_1 \,|\underline{K}_1(\varphi)| + |\underline{K}_1(\varphi)| \cdot \int_1^\infty e^{-s} \cdot s^n \cdot ds \\
&\le c_2 \cdot n! \,|\underline{K}_1(\varphi)|\,.
\end{aligned}$$

In particular, for φ° we may also use Proposition 2 to obtain that

$$\int_{\mathbb{R}^n} e^{-\varphi^\circ} \leq c_2 \cdot n! \cdot \left|\underline{K}_1\left(\varphi^\circ\right)\right| \leq c_2 \cdot 2^n \cdot \left|\underline{K}_1\left(\varphi\right)^\circ\right|.$$

By Proposition 5, $\underline{K}_1(\varphi)$ and $\underline{K}_1(\varphi)^\circ$ are convex bodies. Moreover, since φ is even, $\underline{K}_1(\varphi)$ and $\underline{K}_1(\varphi)^\circ$ are centrally symmetric, and so we may apply the classical Santaló inequality together with the above to obtain that

$$\begin{aligned}\int_{\mathbb{R}^n} e^{-\varphi} \cdot \int_{\mathbb{R}^n} e^{-\varphi^\circ} &\leq c_2^2 \cdot 2^n \cdot n!^2 \left|\underline{K}_1(\varphi)\right| \cdot \left|\underline{K}_1(\varphi)^\circ\right| \\ &\leq c_2^2 \cdot 2^n \cdot n!^2 \left|B_2^n\right|^2 \\ &\leq C^n \cdot n!^2 \left|B_2^n\right|^2\end{aligned}$$

for some $C > 0$. □

4. REVERSE SANTALÓ INEQUALITY FOR THE POLARITY TRANSFORM

Proposition 7. *Let φ be a geometric even convex function on $\mathbb{R}^n$ with $0 < \int e^{-\varphi} < \infty$. Then*

$$0.7 \cdot c^n \cdot \left|B_2^n\right|^2 \leq \int_{\mathbb{R}^n} e^{-\varphi} \cdot \int_{\mathbb{R}^n} e^{-\varphi^\circ}$$

where $c > 0$ is the best constant that can appear in the classical reverse Santaló inequality for convex bodies.

Proof. As in the proof of Proposition 6, we have that $\int_{\mathbb{R}^n} e^{-\varphi} = \int_0^\infty e^{-s} \cdot \underline{K}_s(\varphi)\, ds$. By Proposition 5, $\underline{K}_t(\varphi)$ and $\underline{K}_t(\varphi^\circ)$ are convex bodies (for which the classical reverse Santaló applies) and so by Proposition 2, Cauchy-Schwarz inequality, and the classical reverse Santaló inequality, we obtain that

$$\begin{aligned}\int_{\mathbb{R}^n} e^{-\varphi} \cdot \int_{\mathbb{R}^n} e^{-\varphi^\circ} &= \int_0^\infty e^{-s} \cdot \left|\underline{K}_s(\varphi)\right| ds \cdot \int_0^\infty e^{-u} \cdot \left|\underline{K}_u(\varphi^\circ)\right| du \\ &\geq \int_0^\infty e^{-s} \cdot \left|\underline{K}_s(\varphi)\right| ds \cdot \int_0^\infty e^{-u} \cdot \left|\underline{K}_{1/u}(\varphi)^\circ\right| du \\ &= \int_0^\infty e^{-s} \cdot \left|\underline{K}_s(\varphi)\right| ds \cdot \int_0^\infty \frac{e^{-1/t}}{t^2} \cdot \left|\underline{K}_t(\varphi)^\circ\right| dt \\ &\geq \left(\int_0^\infty \sqrt{e^{-s} \cdot \left|\underline{K}_s(\varphi)\right|} \cdot \sqrt{\frac{e^{-1/s}}{s^2} \cdot \left|\underline{K}_s(\varphi)^\circ\right|} ds\right)^2 \\ &\geq c^n \cdot \left|B_2^n\right|^2 \cdot \left(\int_0^\infty \frac{e^{-s/2-1/(2s)}}{s} ds\right)^2 \\ &\geq a \cdot c^n \cdot \left|B_2^n\right|^2\end{aligned}$$

where

$$a = \left(\int_0^\infty \frac{e^{-s/2-1/(2s)}}{s} ds\right)^2 \geq 0.7$$

may be computed numerically. □

Acknowledgment. The authors would like to thank Keshet Einhorn for her useful comments and corrections.

REFERENCES

1. S. Artstein-Avidan, B. Klartag, and V. Milman, *The Santaló point of a function, and a functional form of the Santaló inequality*, Mathematika **51** (2004), no. 1-2, 33—-48 (2005).
2. S. Artstein-Avidan and V. Milman, *A characterization of the support map*, Adv. Math. **223** (2010), no. 1, 379–391.

3. ———, *Hidden structures in the class of convex functions and a new duality transform*, J. Eur. Math. Soc. (JEMS) **13** (2011), no. 4, 975–1004.
4. K. M. Ball, *PhD dissertation*, Cambridge.
5. J. Bourgain and V. Milman, *New volume ratio properties for convex symmetric bodies in* $\mathbf{R}^n$, Invent. Math. **88** (1987), no. 2, 319–340.
6. M. Fradelizi and M. Meyer, *Some functional forms of Blaschke-Santaló inequality*, Math. Z. **256** (2007), no. 2, 379–395.
7. B. Klartag and V. Milman, *Geometry of log-concave functions and measures*, Geometriae Dedicata **112** (2005), no. 1, 169–182.
8. G. Kuperberg, *From the Mahler conjecture to Gauss linking integrals*, Geom. Funct. Anal. **18** (2008), no. 3, 870–892. MR 2438998 (2009i:52005)
9. J. Lehec, *A direct proof of the functional Santaló inequality*, C. R. Math. Acad. Sci. Paris **347** (2009), no. 1-2, 55–58.
10. ———, *Partitions and functional Santaló inequalities*, Arch. Math. (Basel) **92** (2009), no. 1, 89–94.
11. M. Meyer and A. Pajor, *On the Blaschke-Santaló inequality*, Arch. Math. (Basel) **55** (1990), no. 1, 82–93.
12. V. Milman and L. Rotem, *Mixed integrals and related inequalities*, Journal of Functional Analysis (2012), no. 0, –.
13. F. Nazarov, *The Hörmander proof of the Bourgain-Milman theorem*, Geometric Aspects of Functional Analysis (Mendelson S. Klartag, B. and V.. Milman, eds.), Lecture Notes in Mathematics, vol. 2050, Springer Berlin Heidelberg, 2012, pp. 335–343 (English).
14. R. T. Rockafellar, *Convex analysis*, Princeton Mathematical Series, No. 28, Princeton University Press, Princeton, N.J., 1970.
15. L. Rotem, *Characterization of self-polar convex functions*, Bulletin des Sciences Mathématiques **136** (2012), no. 7, 831 – 838.
16. L. A. Santaló, *An affine invariant for convex bodies of n-dimensional space*, Portugaliae Math. **8** (1949), 155–161.

School of Mathematical Science, Tel-Aviv University, Ramat Aviv, Tel Aviv, 69978, Israel
E-mail address: `shiri@post.tau.ac.il (Artstein-Avidan), boazslom@post.tau.ac.il (Slomka)`